
MONITORING OF SALT TOLERANCE RESPONSES OF DIFFERENT MAIZE INBRED LINES USING PROTEIN PROFILE AND SPECTROFLUORESCENCE.

Nasser H. Abbas, Aisam M. Fayed.

Genetic Engineering and Biotechnology Research Institute (GEBRI), Sadat City University,
Egypt.

Abstract:

Studies were planned to analyze the response of maize genotypes to salt stress (150 mM NaCl) at seedling stage. Six maize inbred lines CML 511, CML 448, CML 444, CML 395, CML 254 and CML 216 were used. The PCR amplification with 8 RAPD primers gave totally 111 RAPD fragments of different molecular weight, in which 77.4 % were polymorphic. The genetic similarities based on Dice's coefficient for each genotype combination were ranged from 0.52 to 0.72. Accessions of CML 511 and CML 448 were included in one main cluster with genetic similarity of 0.65. Leaf proteins were studied by one-dimensional SDS-PAGE. Under non stress conditions, the minor band of 45 kDa was not expressed in varieties of CML 254 and CML 216 while band of 23 kDa was expressed in all accessions except the CML 216. Under salt stress conditions, high molecular weight bands were disappeared and the major protein bands were located at the medium molecular weight range. Protein spectrofluorescence show that inbred lines of CML 511 and CML 448 exhibit closely related fluorescence pattern, this is in context of their genetic similarities as has deduced by RAPD result.

Key words: Maize, Salt stress, RAPD, SDS-PAGE, protein spectrofluorescence.

Introduction:

Salt-affected soils are naturally present in more than 100 countries of the world where many regions are also affected by irrigation induced salinization [1]. The saline soil water inhibits plant growth by an osmotic effect, which reduces the ability of the plant to take up water and by ion excess, which affects the plant cells. Consequently, photosynthetic capacity is reduced and plant senescence is accelerated [2]. Food production is limited by this human-induced salinity, together with the natural and complex salinity found in soils of most semi-arid regions of the world [3].

To avoid this adverse effect, plant evolves a great variety of adaptive mechanisms, such as osmotic adjustment, selective ion uptake, and cytoplasmic and vacuolar ion compartmentation. Recent biotechnical efforts to improve plant salt stress have two main strategies, the elucidation of salt stress signaling and effector output determinants that mediate ion homeostasis. However, the presences of numerous salt tolerance determinants and essential evolutionary necessities have limited the development of salt tolerance transgenic plants [4].

Alternatively, there are crop improvement strategies that are based on the use of molecular marker techniques and biotechnology, and can be used in conjunction with traditional

breeding efforts [5]. RAPD primer showed general consistency as a marker for many plant genera [6]. Genetic variability for salt tolerance was reported in alfalfa [7], Trifolium [8], sunflower [9] and maize [10-12]. In maize, RAPD technique proved to be useful to generate genetic marker especially in cases of inbred lines and haploid lines [13]. The anther molecular marker, stressed protein could be used as important molecular markers for the improvement of salt tolerance using genetic engineering techniques [14]. Proteomics is an increasingly ambiguous term that is now being applied to almost any aspect of protein expression, structure and function. Furthermore, the analysis of the plant's proteome is an important amendment to analysis of the genome, because gene expression is altered under salinity stress [15]. Recently, spectrofluorescent emission of proteins becomes useful tool for protein assay. In this study, we aim to link changes in cellular protein of different maize varieties to their genetic divergence under salt stress condition.

Experimental

Plant growth and treatments

Maize inbred lines, inbred line of 6 maize varieties were used for comparison of salt stress response. These maize varieties have provided by international maize and wheat improvement center (CIMMYT). The pedigree and origin of these varieties are showed in table 1.

Entry	Id. Num.	Acc. Num.	Pedigree	Addresses	Origin of seed	Seed weight	No. of grains	SHU	observations
L1	28110	25835	CML511	A11A0225	TL08A-1903-168	12	50	Cer09-9.141	Lowland
L2	28047	25772	CML448	A10G0632	AF08A-0903-137	13	50	Cert08-13.136	Lowland
L3	28043	25768	CML444	A10G0628	AF08A-0903-134	16	50	Cert08-13.133	Africa MA/ST
L4	25400	22470	CML395	A10G0514	AF08A-0903-121	17	50	Cert08-13.120	Africa MA/ST
L5	17593	16424	CML254	A10G0120	TL08A-1903-185	12	50	Cer09-9.158	Lowland
L6	17123	16394	CML216	A10F0623	AF08A-0903-112	16	50	Cert08-13.111	Africa MA/ST

Table 1. Name, pedigree and origin of inbred lines

Seeds of the maize inbred lines were surface-sterilized for 20 min with 75% ethanol solution, rinsed with sterilized distilled water five times and kept for 4 d at 28°C in sterilized river sand for germination. The seedlings were grown in a greenhouse at 25–31°C under the conditions of a 12 h light period and 60–80% relative humidity. When the seedlings had three full-grown leaves, they were treated by salt stress at 8:00 h by supplementing the nutrient solution with 150 mM NaCl for three successive days. The control seedlings were parallel-treated in the nutrient solution without added NaCl. The stressed and control seedlings were harvested for analysis of gene expression and physiological parameters. The harvested tissue samples were rinsed with sterilized distilled water and frozen in liquid nitrogen.

Apparatus

All fluorescence measurements were carried out on Perkin Elmer L545 fluorescence spectrometer in the range (260-750 nm). SDS-PAGE electrophoresis has been carried on a Mini-Protein II TM Cell (Bio-Rad Richmond, CA, USA). Amplification was performed in a Thermal Reactor (TPersonal, Biometra).

General procedure

The genetic characterization and genetic diversity assessment were conducted using protein and RAPD markers. The extraction of proteins was performed using 0.2 g of leaves as described by [16]. Protein crude extract was used for assays and quantification as described by [17] using bovine serum albumin as a standard. Proteins were separated by 12% acrylamide gel electrophoresis (SDS-PAGE) using equal (30 µg) protein content and stained by Coomassie Brilliant Blue (Sigma Aldrich) according [18]. For spectrofluorescence assay protein has dialyzed against 10% glycerol and 50 mM HEPES, pH 7.0 for 18 hrs. The genomic DNA was isolated from leaf tissue following the CTAB procedure described by [19]. RAPD-PCR amplification was performed with 8 RAPD primers as shown in (Table 2). The PCR programme had 36 cycles in which first denaturation was carried out at 94°C for 3 min, segment denaturation at 94°C for 1 min, annealing at 35.5°C for 1 min, extension at 72°C for 2 min and final extension for 3 min at 72°C. RAPD fragments were separated on 1.4. % agarose gels in 0.5x TBE buffer. Presense/absence of protein fractions or DNA fragments was transformed to binary data (0 and 1). The computing of binary data including coefficients of similarity and UPGMA clustering was performed using NTSYS-pc software [20].

5'-3' sequence	patterns RAPD primers	5'-3' sequence	patterns RAPD primers
<i>GGACCCAACC</i>	<i>OPD-02</i>	<i>ACGCACAACC</i>	<i>OPE-15</i>
<i>CCTTGACGCA</i>	<i>OPB-12</i>	<i>CACTCTCCTC</i>	<i>OPH-17</i>
<i>GGTACTCCCC</i>	<i>OPN-03</i>	<i>ACCGCGAAGG</i>	<i>OPD-01</i>
<i>AGTCAGCCAC</i>	<i>OPA-03</i>	<i>ACAAC TGGGG</i>	<i>OPN-10</i>

Table2: RAPD primers used in this study which gave clear and reproducible bending

Fluorescence emission measurements

Measurement of leaf water soluble protein fluorescence was determined with Perkin Elmer L545 fluorescence spectrometer. The excitation wavelength was 270 nm. The protein sample was suspended in 200 µl of 10% (v/v) glycerol, 50 mM HEPES, pH 7.0.

RESULTS

RAPD Data Analysis

PCR amplification of genomic DNA was tested on 13 RAPD primers in two rounds of amplification which of 8 primers gave clear and reproducible banding patterns (Fig. 1). PCR amplification with 8 RAPD primers gave totally 111 RAPD fragments of different molecular weight, which of 77.4 % were polymorphic (in presence/absence of bands). The Similarity coefficients were calculated according to dice matrix [21]. The genetic coefficient distance for each genotype combination ranged from 0.52 to 0.72. The similarity dendrogram in (fig. 2) shows that the accessions can be classified into two major groups A and B. The first main cluster (A) contains L1 and L2 accessions that are separated at genetic similarity of 0.65. The second main clusters (B) separated at genetic similarity 0.62 into two main sub clusters. The first main sub cluster (1B) separated into two small sub clusters, 1Ba sub cluster included L3 and L4 at genetic similarity of 0.72 and the 1Bb sub cluster included individual cultivars L5 at genetic similarity of 0.67. The second main sub cluster (2B) included individual cultivars L6 at genetic similarity of 0.62.

Protein profile:

The protein banding pattern of non-salt treated and salt treated maize leaf is demonstrated in (Fig. 3). The six lanes represent six maize varieties total cellular protein patterns. The protein profile of non-treated leaf show major protein bands pattern of 320 kDa, 280 kDa, 220 kDa, 55 kDa, 41 kDa and 36 kDa. Under non salt stress conditions, the six maize varieties show no difference among major protein bands however, minor protein bands of 45 kDa and 23 kDa are absent in protein profile of L6 maize variety. Under salt stress conditions, the protein banding pattern of maize leaf varieties show absence of high molecular protein. The major protein bands of salt stress conditions are 55 kDa, 45 kDa, 34 kDa, 28 kDa, 24 kDa and 23kDa. The number of bands of leaf protein extracted from control plants was much higher than of the salt treated plants (Fig. 3).

Spectroscopic analysis:

The emission fluorescence of extracted leaf protein samples of control and salt stressed were measured using visible and UV wave length. The emission fluorescence of all sample were recorded at excitation spectra of 270 nm. All maize varieties show fluorescence emission peak at range of 320 to 340 nm however, the intensity of this peak show alternation of intensity for salt stressed and control samples. Maize varieties L1 and L2 have higher peak intensity of salt stressed leaf protein compared to the control samples. The other maize varieties L3, L4, L5, L6 shows higher peak intensities for the control samples. Other emission peaks were associated

with salt stressed protein of samples varieties L2, L3, L4 and 6 with two common peaks at 440 and 490 nm. The maize variety number 6 shows excess emission peak at 550 nm for the control sample.

DISCUSSION

RAPD assay can efficiently generate both randomly dispersed markers as well as markers linked to specific genes. Constraint of the technique is its reproducibility, which difficult possibility of interchanging results in and between laboratories. However, the problems related with reproducibility of RAPD can be resolved by rigorous attention to detail, [22]. In this study, the RAPD analysis with 8 random primers gave totally 111 fragments which of 77.4 % were polymorphic. These results are close with results of study of 57 elite corn inbred lines where 84% RAPD fragments were polymorphic [23]. Other result on hybrid lines of maize show similar ratio of polymorphic bands, [24]. The genetic similarities dendrogram shows that the accessions from the same geographical region were found to have a close genetic relationship (Fig. 2).

The molecular mechanism of salt stress tolerance in plant could be elucidated using cDNA microarrays. However, mRNAs may not be transcribed or that changes in the protein level or enzyme activity can occur without any detectable change in transcript abundance due to translational or other levels of control. Therefore, it is necessary to study the salt stress responses at the protein level [25]. In this study the major protein profile of maize leaf under control conditions show protein bands that can categorized in high molecular weight (estimated as 320, 280 and 220 kDa), medium range molecular weight (55, 41 and 36 kDa) and low range molecular weight less than 25 kDa. The high molecular weight bands in maize leaf are not recognized in previous research of maize [26]. This could be because protein profile of plant leaves is significantly influenced by the extraction method [27].

The major band of 55 kDa is closely related to the major mesophyll protein band of 56 kDa of ribulose-1,5-bisphosphate carboxylase/oxygenase in maize. This chloroplast expressed enzyme is crucial for the fixation of carbon dioxide [28,29]. Many of maize cytosolic cellular protein in maize have molecular weight ranged 45-29 kDa. Most of these proteins are enzymes involved in glycolytic pathway and are necessary for cellular energy conservation [30].

Under non salt stress different maize varieties show some specific variation. Minor band of 45 kDa did not expressed in varieties of L5 and L6 while another minor band of 23 kDa was not expressed in L6 variety. The differential pattern of salt stress protein is highly related to the plant genotype. Consequently, some plants did not display large differences in protein patterns between control and salt stressed conditions, in other cases the stressed protein SDS PAGE profile show a contrasted profile versus unstressed protein profile [31,32]. The protein profile patterns emphasize the closely genetic relationship between L5 and L6 variety. Protein bands within the range of 20–45 kDa were isolated from the apoplast infiltrate of maize leaf [33]. Under salt stress conditions, protein profile of the maize varieties show high homogeneity pattern among the different varieties. Comparing to non-stress protein pattern, the high molecular weight bands were disappeared and the major protein bands were located at the medium molecular weight range. This could be due to the inhibitory effects of salt stress on transcriptional process [34]. The fast breakdown of cellular protein during salt stress might be also due to the increasing activity of acid and alkaline protease [35]. Homeostasis and mineral deficiencies has been a consequence of salt stress [36]. In maize, mineral deficiency leads to increase in chloroplast

protein content. The identified maize chloroplast protein show 17.5 kDa for coupling factor, 23 kDa for structure protein, 33 kDa for cytochrome F, 37 kDa coupling factor, 42 kDa for cytochrome b6, 45 kDa for transhydrogenase and 56 kDa for coupling factor and ribulose diphosphate carboxylase (Barrand and Crane, 1974). Several proteins within the range of 56 kDa-17 kDa have been identified as apoplast proteins for biotic stress in maize [37].

The fluorescence emission of proteins is mainly referred to their contents of aromatic amino acids tryptophan, tyrosine and phenylalanine [38]. The main peak of fluorescent emission in all fluorescent patterns of normal and salt stress conditions were at 320 nm except variety L1 which was at 340 nm (Fig. 4). This fluorescent emission value is very close to the emission peak of tryptophan amino acid [39,40]. The tryptophan residue in protein has emission spectrum a ranged from 310 ~ 350 nm should reflect the average environment of the tryptophan. Hence, the variations in tryptophan emission are due to the structure of the protein [41]. The increase of emission wave length of tryptophan peak in sample 2 could refer as a change in its protein structure comparing to other maize varieties. The fluctuation pattern of peak intensity between salt and control protein samples show higher peak intensity in case of control samples (with exception of varieties L1 and L2). The quenching effect in case of salt stressed samples could be result of protein binding interaction that cause collisional quenching of emission intensity [42]. The increment of emission spectra that associate salt stressed protein samples of varieties L1 and L2 indicate protein changes that prevent quenching effect of protein collision. The other important observation is in case of losing the quenching effect the salt stress protein samples of varieties L1 and L2 show higher peak intensity rather than the average intensity of control samples. Several data show increase of protein fluorescence intensity as a result of protein phosphorylation process or protein binding with calcium [43,44]. Under salt stress, plants activate sensor proteins through protein binding with calcium and phosphorylation of many protein kinases [45]. Our data suggest presence of protein phosphorylation and or protein calcium binding in case of salt stressed protein. However, the subsequent effect of emission intensity increment was hidden by the collisional protein effect. Additional peaks of 380 nm, 420 nm and 440 nm were associated with salt stressed samples of varieties L2, L3, L4 and L6. These emission spectra are most likely due to emission spectra of non-pretentious compounds that may co-precipitate during protein preparation. Some maize secondary metabolites show fluorescence spectra close to peaks in our results. Gallic acid, o-Coumaric acid and Cinnamic acid have emission peaks of 382 nm, 426 nm and 420 nm respectively. These substances have antioxidant effect and research data show their involvement in abiotic resistance of maize plant [46-48]. These data may interpret the presence of 380 nm, 420 nm, and 440 nm in most spectrofluorescence samples of salt stress under study.

CONCLUSION:

According RAPD data the maize varieties under study can grouped into two main groups where varieties L1 and L2 has grouped in one cluster. Results of SDS-PAGE and protein spectrofluorescence show that, the main differences between maize varieties have observed under control condition. However, protein spectrofluorescence data show similar behavior of L1 and L2 varieties which are in context with result of RAPD genetic similarities. Protein spectrofluorescence has proved to be a reliable tool for monitoring proteome responses to salt stress condition.

AKNOWLEDGMENT

I would thank Dr. Kamal Abdulatif (bioinformatics department, GBRI) for his helping of RAPD result analysis.

Primer name	Total No. bands	Polymorphic bands	% of Polymorphism
OPA03	21	16	76
OPA07	19	19	100
OPD01	13	8	61.5
OPE15	9	8	88.9
OPH17	7	7	100
OPN03	13	13	100
OPN10	14	11	78.5
OPP12	15	12	80

Table 3. Total band number and polymorphic Bands of eight RAPD primers used with the 6 maize inbred lines




Fig.1. The random amplified polymorphic DNA (RAPD) profile of maize varieties using the random primers which is listed in Table 2




Fig. 2. RAPD based dendrogram of investigated maize genotypes based on cluster analysis of Nei and Li's genetic distance (1979).




Fig. 3. SDS-PAGE profiles of total leaf protein of maize varieties under control condition (A) and under salt stress condition (B).




Fig.4. Fluorescence emissionspectra (excitation: 270 nm) of total leaf protein of the six maize varieties in water. Leaf maize protein under control condition (N) and salt stress condition.

REFERENCES

- 1- **Hillel, D., (2005).** Soil salinity: Historical and contemporary perspectives. *Proceedings of the international salinity forum*, Riverside, 235–240.
- 2- **Jones, J. B., (2003).** Agronomic Handbook: Management of Crops, Soils and Their Fertility. *CRC Press. Boca Raton*, p. 450.
- 3- **Rengasamy, P., (2010).** Soil processes affecting crop production in salt-affected soils. *Functional Plant Biology*, **37**, 613–620.
- 4- **Munns, R., (2005).** Genes and salt tolerance: bringing them together. *New Phytologist*, **167**, 645–663.
- 5- **Ribaut, J. M. and Hoisington, D., (1998).** Marker-assisted selection: new tools and strategies. *Trends in Plant Sci.*, **3**, 236-239.
- 6- **Fritsch, P., Hanson, A. M., Spore, D. C., Pack, E. P. and Riesberg, H. L., (1993)** Constancy of RAPD primer amplification strength among distantly related taxa of flowering plants. *Plant Molecular Biology Reporter*, **11**, 10-20.
- 7- **Mckimmie, T. and Dobrenz, A. K., (1991).** Ionic concentrations and water relations of alfalfa seedlings differing in salt tolerance. *Agron. J.*, **83**: 363-367.
- 8- **Ashraf, M., McNelly, T. and Bradshaw, A. D., (1987).** A developmental window for salt adaptation in Sorghum bicolor. *J. Exp. Bot.*, **44**, 645-652.
- 9- **Francois, L. E., (1996).** Salinity effects on four sunflower hybrids. *Agron. J.*, **88**: 215-219.
- 10- **Maiti, R. K., Amaya, L. E., Cardona, S. I., Dimas, A. M., De La Rosa-Ibarra, M. and Castillo, H. D., (1996).** Genotypic variability in maize cultivars (*Zea mays* L.) for resistance to drought and salinity. *J. Plant Physiol.*, **148**, 741- 744.
- 11- **Sharif, A., Rasul, E., Nisar, A. and Sadiq, M., (1999).** Response of maize (*Zea mays* L.) genotypes to NaCl salinity at various growth stages of crop. *Pak. J. Biol. Sci.*, **23**, 606-608.
- 12- **Khan, A. A., Sajjad, A. R. and McNeilly, T., (2003).** Assessment of salinity tolerance based upon seedling root growth response functions in maize (*Zea mays* L.). *Euphytica*, **131**: 81-89.
- 13- **Beaumont, H. V., Mantet, J., Rocheford, R. T. and Widholm, M. J., (1996).** Comparison of RAPD and RFLP markers for mapping F2 generations in maize (*Zea mays* L.). *Theoretical and Applied Genetics*. **93**, 606-612.
- 14- **Pareek, A., Singla, S. L. and Grover, A., (1997).** Salt Responsive Proteins/Genes in Crop Plants, Strategies for Improving Salt Tolerance in Higher Plants. New Delhi, Oxford and IBH, 365–391.
- 15- **Zörb, C. S., Schubert, A., Neeb, D., Karl, M., Linder and S. Schubert, (2004).** The biochemical reaction of maize (*Zea mays* L.) to salt stress is characterized by a mitigation of symptoms and not by a specific adaptation. *Plant Sci.*, **167**, 91–100.
- 16- **Wang, W., Scali M., Vignani, R., Spadafora, A., Sensi, E., Mazzuca, S. and Cresti, M., (2003).** Protein extraction for two-dimensional electrophoresis from olive leaf, a plant tissue containing high levels of interfering compounds. *Electrophoresis*, **24**, 2369–2375.
- 17- **Bradford, M. M., (1976).** A Rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, **72**:248-254.
- 18- **Hames, B. D. and Rickwood, D., (1990).** Gel Electrophoresis of Proteins. *A Practical Approach*, New York: Oxford Univ. Press.

- 19-Porebski S, Bailey LG, Baum BR (1997)** Modification of a CTAB DNA extraction protocol for plants containing high polysaccharides and polyphenol component. *Plant Molecular Biology Reporter* **15**, 8-15.
- 20-Rohlf, F. J., (2000).** NTSYS-pc: numerical taxonomy and multivariate analysis system, version 2.1. Exeter Software: Setauket, NY.
- 21-Nei, M. and Li, W.H., (1979).** Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci.*, **76**, 5269- 5273.
- 22-Jones, C. J., Edwards, K. J., Castiglione, S., Winfield, M. O., Sala, F., Van-der-Weil, C., Vosman, B. L., Matthes, M., Daly, A., Brettschneider, R., Bettini, P., Buiatti, M., Maestri, E., Marmioli, N., Aert, R. L., Volckaert, G., Rueda, J., Vazquez, A., Karp, A., (1998).** Reproducibility testing of RAPDs by a network of European laboratories. In: Karp A, Isaac PG, Ingram DS (eds.) *Molecular tools for screening biodiversity*, Chapman & Hall, London, UK, pp 176-179.
- 23-Hahn, V., Blankenhorn, K., Schawall , M. and Melchinger, A. E., (1995)** Relationships among early European maize inbreds: III. Genetic diversity revealed with RAPD markers and comparison with RFLP and pedigree data. *Maydica* 40:299-310. BAUER, I., S. MLADENOVIC DRINIC, M. FILIPOVIC, and K. KONSTANTINOV (2005): Genetic characterization of early maturing maize hybrids (*Zea mays* L.) obtained by protein and RAPD markers. *Genetika*, **37**, 235.
- 24-Bauer, I., Mladenovic, S. D., Filipovic M., and Konstantinov, K., (2005)** Genetic characterization of early maturing maize hybrids (*Zea mays* L.) obtained by protein and RAPD markers. *Genetika*, **37**, 235.
- 25-Shunping, Y., Tang, Z., Su, W. and Sun, W., (2005).** Proteomic Analysis of Salt Stress-Responsive Proteins in Rice Root. *Proteomics*, **5**, 235–244.
- 26-Mohamed, A. A., (2005).** Two-dimensional Electrophoresis of Soluble Proteins and Profile of Some Isozymes Isolated from Maize Plant in Response to NaCl. *Research Journal of Agriculture and Biological Sciences*, **1**, 38-44.
- 27-Maldonado , A. M., Echevarria-Zomeno, S., Jean-Baptiste, S., Hernandez, M., Jorri-
Novo, J. V., (2008).** Evaluation of three different protocols of protein extraction for *Arabidopsis thaliana* leaf proteome analysis by two-dimensional electrophoresis. *Journal of Proteomics*, **71**, 461-472.
- 28-Klein, S. M., and Vernon, P. L., (1973).** Protein composition of spinach chloroplasts and their photosystem I and photosystem II fragments. *Photochem. Photobiol.* Bewley, J.D. and Black, M. (1994) *Seed: Physiology of Development and Germination*. Plenum Press.
- 29-Bewley, J. D. and Black, M., (1994).** *Seed: Physiology of Development and Germination.* Plenum Press, New York.
- 30-Okamoto, T., Higuchi, K., Shinkawa, T., Isobe, T., Lörz, H., Koshiba, T. and Kranz, E., (2004).** Identification of Major Proteins in Maize Egg Cells. *Plant Cell Physiol.* , **45**, 1406–1412.
- 31-Amini, F., Ehsanpour, A. A., Hoang, T. Q. and Shin Sh. J., (2007).** Protein Pattern Changes in Tomato under In Vitro Salt Stress. *Russian Journal of Plant Physiology*, **54**, No. 4, pp. 464–471
- 32-Sohrabi, N., Tajabadipour, A., Motamed, N. and Seyedi, M., (2011).** A Change in Leaves Protein Pattern of Some Pistachio Cultivars under Salinity Condition. *International Journal of Nuts and Related Sciences*, **2**, 67-74.

- 33- Witzel, K., Shahzad, M., Matros, A., Mock, P. H. and Mühling, H. K., (2011).** Comparative evaluation of extraction methods for apoplastic proteins from maize leaves. *Plant Methods*, **7**, 48-59.
- 34- Jain, S., Nainawatee, H. S., Jain, R. K. and Chowdhury, J. B., (1993).** Salt-tolerance in Brassica juncea L. II. Salt-stress induced changes in polypeptide pattern of in vitro selected NaCl-tolerant plants. *Euphytica.*, **65**, 107-112.
- 35- Parida, A., Das, A. B., Mitra, B. and Mohanty, P., (2004).** Salt stress induced alterations in protein profile and protease activity in the Mangrove Bruguiera Parviflora. *Z. Naturforsch.*, **59**, 408- 414.
- 36- Schroepel-Meyer, G. and Kaiser, W. M., (1988).** Ion osmeostasis in chloroplast under salinity and mineral deficiency. I. Solute concentrations in leaves and chloroplast from spinach plants under NaCl and NaNO₃ salinity. *Plant Physiology*, **87**, 822–827.
- 37- Teale, W. J. F., (1960).**The ultraviolet fluorescence of protein in neutral solution. *Biochem.J.*, **76**, 381-388.
- 38- Boteva, R., Zlateva, T., Dorovska-Taran, V., Visser, A. J., Tsanev, R., Salvato B., (1996).** Dissociation equilibrium of human recombinant interferon. *Biochemistry* **35**,14825–14830.
- 39- Kabiri, M., Amiri-Tehranizadeh, Z., Baratian, A., Saberi, R. M. and Chamani, J., (2012).** Use of Spectroscopic, Zeta Potential and Molecular Dynamic Techniques to Study the Interaction between Human Holo-Transferrin and Two Antagonist Drugs: Comparison of Binary and Ternary Systems. *Molecules*, **17**, 3114-3147.
- 40- Vivian, J. T. and Callis, P. (2001).**Mechanisms of tryptophan fluorescence shifts in proteins. *Biophys J.*, **80**, 2093–2109.
- 41- Eftink, M. R., (1991).** Fluorescence quenching: theory and applications. *Topics in fluorescence spectroscopy*, **2**: 53–126.
- 42- Miranda, F. F., Thorolfsson, M., Teigen, K., Sanchez-Ruiz, M. J. AND Martinez, A., (2004).** Structural and stability effects of phosphorylation: Localized structural changes in phenylalanine hydroxylase. *Protein Science*, **13**,1219–1226.
- 43- VanScyoc, W. S., Sorensen, B. R., Rusinova, E., Laws, W. R., Ross, J. B. and Shea M. A., (2002).** Calcium binding to calmodulin mutants monitored by domain–specific intrinsic phenylalanine and tyrosine fluorescence. *Biophys J.*, **83**, 2767-2780.
- 44- Du, W., Lin, H., Chen, S., Wu, Y., Zhang, J., Fuglsang, T. A., Palmgren, G. M., Wu, W. and Guo, Y., (2011).** Phosphorylation of SOS3-Like calcium-binding proteins by their interacting SOS2-like protein kinases is a common regulatory mechanism in arabidopsis. *Plant Physiology*, **156**, 2235–2243.
- 45- Lopez-Martinez, X. L., Oliart-Ros, M. R., Valerio-Alfaro, G., Lee, C., Parkin, L. K. and Garcia, S. H., (2009).** Antioxidant activity, phenolic compounds and anthocyanins content of eighteen strains of Mexican maize. *Food Science and Technology*, **42**, 1187–1192.
- 46- Pál, M., Horváth, E., Janda, T., Páldi, E. and Szalai, G., (2005).** Cadmium stimulates the accumulation of salicylic acid and its putative precursors in maize (*Zea mays*) plants. *Physiologia Plantarum*, **125**, 356–364.
- 47- Tuberoso, G. C., Kowalczyk, A., Sarritzu, E. and Cabras, P., (2007).**Determination of antioxidant compounds and antioxidant activity in commercial oilseeds for food use. *Food Chemistry*, **103**, 1494–1501.